

eISSN: 09748369, www.biolmedonline.com

Cytotoxic effect of Carvacrol on human cervical cancer cells

#Mehdi SJ¹, #Ahmad A², Irshad M¹, Manzoor N², *Rizvi MMA¹¹ Genome Biology Lab, Department of Biosciences, Jamia Millia Islamia, New Delhi-25, India.² Enzyme Kinetics Lab, Department of Biosciences, Jamia Millia Islamia, New Delhi-25, India.

Both the authors have equal contribution

*Corresponding Author: rizvijmi@gmail.com

Abstract

Carvacrol is a component of numerous aromatic plants which has been evaluated for substantial pharmacological properties. Although the carvacrol induced cytotoxicity has already been reported, but no such study have been made on human cervical cancer, HeLa and SiHa cells. Therefore, in the present study an attempt has been made to investigate the cytotoxic effect of carvacrol on cervical cancer cells. Cytotoxicity induced by carvacrol was determined by different assays like MTT assay and LDH assay. Apoptosis was measured by DNA fragmentation assay. The study clearly showed the dose dependent cytotoxic effect of carvacrol in HeLa and SiHa cells at an IC₅₀ of 50 mg L⁻¹ by both the cytotoxic assays respectively. The dying cells showed characteristics of apoptosis such as, DNA fragmentation. The data in the present study clearly demonstrated cytotoxic effects of carvacrol on human cervical cancer cells. Carvacrol could have a potential therapeutic significance in treating cancer.

Keywords: Carvacrol; Cervical cancer; HeLa cells; SiHa cells; Apoptosis.

Introduction

In recent years, the interest in medicinal plants and their biologically active derivatives has increased, in relation to the possible development of novel potential drugs for several pathologies of relevant social impact (Hedberg, 1993; Heinrich and Gibbons, 2001). It is well known that natural products from the extracts of medicinal plants are used in the treatment of skin, respiratory, neuromuscular and mental health disorders and also in obstetrics and gynecology (Abo *et al.*, 2000; Ahmad *et al.*, 1998; Ankli *et al.*, 2002; Dutta *et al.*, 1998; Pinn, 2001). The anti-tumor activity and the possible applications of medicines from medicinal plants for cancer prevention have been recently described (Chiu and Wu, 2002; Cragg and Newman, 1999; Katsube *et al.*, 2003; Mukherjee *et al.*, 2001; Richardson, 2001; Tatman and Mo, 2002; Wargovich *et al.*, 2001; Popov *et al.*, 2001; Ruffa *et al.*, 2001).

Essential oils and their components have been timely honored for their pharmaceutical properties. Plant-derived substances have recently become of great interest owing to their versatile applications (Baris *et al.*, 2006; Khan *et al.*, 2010). The development of pharmaceuticals begins with identification of active principles, detailed biological assays and dosage formulations, followed by clinical studies to establish safety, efficacy and pharmacokinetic profile of new drug

(Iwu *et al.*, 1999). Carvacrol (5-isopropyl-2-methylphenol) (Figure 1) is a natural isopropyl cresol and is credited with a series of pharmacological properties including antimicrobial and antifungal effects (Pina-Vaz *et al.*, 2004; Braga *et al.*, 2007; Chami *et al.*, 2005). Carvacrol is generally recognized as a safe food additive and used as a flavoring agent in baked foods, sweets, beverages and chewing gum (Fenaroli, 2002). It is also well known that essential oils, which are rich in carvacrol, possess strong antioxidant properties equivalent to those of ascorbic acid, butyl hydroxyl toluene (BHT) and vitamin E (Aeschbach *et al.*, 1994; Mastelic *et al.*, 2008). Since many antioxidants exert anticarcinogenic effects (Kallistratos *et al.*, 1994; Evangelou *et al.*, 1997; Liasko *et al.*, 1998; Karkabounas *et al.*, 2002), it is possible that carvacrol functions in a similar way and the anti-proliferative properties of carvacrol on non-small cell lung cancer cells, A549, chronic myeloid leukemia cells, K562, murine B16 melanoma cells and human metastatic breast cancer cell line, MDA-MB 231 have been shown (He *et al.*, 1997; Horvathova *et al.*, 2007; Karkabounas *et al.*, 2006; Koparal and Zeytinoglu, 2003; Lampronti *et al.*, 2006; Arunasree, 2010).

Based on this hypothesis and reported data, the present study is aimed to evaluate the cytotoxic effects of carvacrol on human cervical cancer HeLa cells and SiHa cells. Quercetin at

an IC_{50} value of 50 μM was used as positive control to assess the efficacy of carvacrol (Wei *et al.*, 2007).




Figure I: Chemical structure of Carvacrol.

Materials and Methods

Chemicals

Carvacrol (98%) and Quercetin ($\geq 95\%$) (quercetin was used as positive control) were purchased from Sigma–Aldrich (USA). DMEM (Dulbecco's modified Eagle's medium), MTT (3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl tetrazolium bromide) and 0.25% trypsin and 0.02% EDTA mixture was purchased from Himedia (India). Cytoscan™-LDH Cytotoxicity Assay Kit was purchased from GBiosciences (USA). Fetal bovine serum (FBS) was from Biowest (USA). All the other chemicals and reagents were purchased from local companies and were of molecular biology grade. Carvacrol and Quercetin were dissolved in DMSO and stored at 4°C.


Cell lines and culture conditions

Human cervical cancer cell lines, HeLa and SiHa were obtained from National Centre for Cell Sciences (NCCS) Pune, India. Both the cell lines were grown as monolayer cultures in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum and antibiotics (100 units/ml penicillin and 100 $mg L^{-1}$ streptomycin) in a humidified atmosphere of 5% CO_2 at 37°C in T-75 flasks and were sub cultured twice a week.

Cytotoxicity assay

The cytotoxic effect of carvacrol was assessed in human cervical cancer HeLa and SiHa cells by the MTT assay (Chou and Talalay, 1984). Briefly, cells were seeded at a number of 2×10^4 per well onto 96-well plates (200 μl /well) in triplicates, allowed to attach and grow for 24 h and subsequently exposed to different carvacrol ($25 mg L^{-1}$ - $500 mg L^{-1}$) concentrations for 48 h. At the end of the treatment, the medium was removed and cells were incubated with 20 μl of MTT (5 mg/ml in PBS) in fresh medium for 4 h at 37°C. After four hours, formazan crystals, formed by mitochondrial reduction of MTT, were solubilized in DMSO (150 μl /well) and the absorbance was read at 570 nm after 10 min incubation on the iMark Microplate Reader (BioRad, USA). Percent inhibition of cytotoxicity was calculated as a fraction of control (without carvacrol) and the cytotoxicity of carvacrol was expressed as IC_{50} .

Figure II: Dose-dependent effect of carvacrol on SiHa and HeLa cell proliferation. Both the cells were cultured in 10% FBS medium and treated with 25, 50, 100, 200 and 500 $mg L^{-1}$ carvacrol for 48 h and cell proliferation was monitored by MTT assay. The percent viable cells were calculated in comparison to untreated cells taken as 100%. Values were expressed as mean \pm SD and the experiment was performed in triplicate ($P < 0.05$).



LDH assay

Cytotoxicity induced by carvacrol was assessed by lactate dehydrogenase (LDH) leakage into the culture medium. Following exposure to the carvacrol the culture medium was aspirated and centrifuged at 3000 rpm for 5 min in order to obtain a cell free supernatant. The activity of LDH in the medium was determined using a commercially available Cytoscan™-LDH Cytotoxicity Assay Kit. Percent inhibition of cytotoxicity was calculated as a fraction of control (without carvacrol) and the cytotoxicity of carvacrol was expressed as IC₅₀.

Detection of apoptotic DNA fragments

DNA fragmentation was detected by agarose gel electrophoresis. 1X10⁶ HeLa and SiHa cells were plated in 30 mm culture plate. When the cells reached approximately 70% confluency, increasing concentrations of carvacrol (25 mg L⁻¹-500 mg L⁻¹) were added and the cells were incubated for 48 h. After 48 hours, cells were harvested and pelleted by centrifugation (Eppendorf 5804R, Germany). Cellular DNA was extracted by SDS/proteinase K treatment, phenol-chloroform extraction, and ethanol precipitation as described previously (Sambrook and Russell, 2001) and then dissolved and stored in TE buffer and the DNA samples obtained were analyzed by 2% agarose gel electrophoresis. After electrophoresis, the gels were stained with ethidium bromide, and visualized as a DNA ladder with UV.

Statistical analysis

Viable cell ratios were evaluated by Student 't' test. The data were presented as arithmetic mean ± standard deviation. A value of P < 0.05 was considered to be statistically significant.

Results

Carvacrol inhibited proliferation of HeLa and SiHa cells

Data on the cytotoxic effects of carvacrol using two human tumor cell lines *in vitro* are showed in Figure II and III. Carvacrol has cytotoxic effects *in vitro* at clinical acceptable concentrations (IC₅₀ values ≤ 50 mg L⁻¹) by MTT methods and LDH methods respectively. The cytotoxic effect of carvacrol was determined using concentrations ranging 25 mg L⁻¹- 500 mg L⁻¹ for 48 h. After 48 h exposure, carvacrol induced concentration-dependent cytotoxic effects in cervical cell lines with IC₅₀ of 50±3.89 mg L⁻¹ and 50±5.95 mg L⁻¹ in SiHa and HeLa cells, respectively using MTT method and 55±0.15 mg L⁻¹ and 50±0.05 mg L⁻¹ in SiHa and HeLa cells, respectively by LDH method.

Validation of apoptosis measurement by DNA laddering

In our results, the cells were treated with carvacrol, and the DNA was directly extracted and run on agarose gel. DNA hyperfragmentation, if presented, was seen as a stepwise ladder of DNA fragments.

Figure III: Dose-dependent effect of carvacrol on SiHa and HeLa cell proliferation. Both the cells were cultured in 10% FBS medium and treated with 25, 50, 100, 200 and 500 mg L⁻¹ carvacrol for 48 h and cytotoxicity was monitored by LDH assay. The percent cytotoxicity was calculated in comparison to untreated cells taken as 100%. Values were expressed as mean ± SD and the experiment was performed in triplicate (P<0.05).





Figure IV: Analysis of DNA fragmentation in human cervical cancer cells treated with carvacrol. **a.** HeLa cells treated with carvacrol. Lane 1: 100 bp DNA ladder; lane 2: Control cells; lane 3: Cells treated with 25 mg L⁻¹ carvacrol; lane 4: Cells treated with 50 mg L⁻¹ carvacrol; lane 5: Cells treated with 100 mg L⁻¹ carvacrol; lane 6: Cells treated with 200 mg L⁻¹ carvacrol; lane 7: Cells treated with 500 mg L⁻¹ carvacrol; lane 8: Cells treated with 50 μM quercetin. **b.** SiHa cells treated with carvacrol. Lane 1: 100 bp DNA ladder; lane 2: Control cells; lane 3: Cells treated with 25 mg L⁻¹ carvacrol; lane 4: Cells treated with 50 mg L⁻¹ carvacrol; lane 5: Cells treated with 100 mg L⁻¹ carvacrol; lane 6: Cells treated with 200 mg L⁻¹ carvacrol; lane 7: Cells treated with 500 mg L⁻¹ carvacrol; lane 8: Cells treated with 50 μM quercetin.



The data (Figure IV) shows that DNA laddering is pronounced for carvacrol (50 mg L⁻¹) in HeLa cells and (50 mg L⁻¹) in SiHa cells respectively. These results confirm that carvacrol can induce apoptosis of HeLa and SiHa cells.

Discussion

Research on biologically active compounds from essential oils has proved them to be potent antibacterial, antifungal and anti-oxidant agents (Lampronti *et al.*, 2006; Albuquerque *et al.*, 2007; Ao *et al.*, 2008; Baik *et al.*, 2008; Bakkali *et al.*, 2008). Carvacrol (5-isopropyl-2-methylphenol) is one of main substances of essential oil possess antimicrobial and antifungal effects (Pina-Vaz *et al.*, 2004; Braga, 2007; Chami *et al.*, 2005).

Cellular proliferation depends on the rates of cell division and death and, thus, many anticancer drugs have been used to prevent cancer cell division in order to inhibit cancer cell proliferation. In vitro cytotoxicity assays can be used to predict human toxicity and for the general screening of chemicals (Clemedson and Ekwall, 1999; Scheers *et al.*, 2001). It has been previously reported that different cytotoxicity assays can give different results depending on the test agent used and the cytotoxicity assay employed (Weyermann *et al.*, 2005). The results obtained from the cytotoxicity assays indicate that inhibition of HeLa and SiHa cells was

gradually increased by the addition of carvacrol in MTT and LDH assays. It reveals that HeLa and SiHa cell line is susceptible to carvacrol. Although, the LDH leakage assay is based on the release of the enzyme into the culture medium after cell membrane damage whereas the MTT assay is mainly based on the enzymatic conversion of MTT in the mitochondria, both the cytotoxicity assays, employed to assess carvacrol toxicity *in vitro*, showed the similar results.

Apoptosis is a physiological process of cell elimination, and DNA fragmentation is one of the hallmarks of cell apoptosis. The results showed the apoptosis proportion of cells was increased by treatment of carvacrol in both the cervical cancer cell lines.

Conclusion

In summary, the present study demonstrated that carvacrol is a potent anti-cancer compound with an IC₅₀ of 50 mg L⁻¹ at 48 h inducing growth inhibition in both the human cervical cancer cells. Further research based on animal models may resolve *in vivo* efficacy of carvacrol.

Acknowledgement

The financial assistance provided by the Defense Research and Development Organization (DRDO), New Delhi, is gratefully acknowledged. The assistance of Mr. Irfan Ahmad is acknowledged with thanks.

References

- Abo KA, Adeyemi AA, Adeite DA, 2000. Ethnobotanical survey of plants used in the treatment of infertility and sexually transmitted diseases in southwest Nigeria. *African Journal of Medicine and Medical Sciences*, 29: 325–327.
- Aeschbach R, Loliger J, Scott BC, Murcia A, Butler J, Halliwell B, Aruoma OI, 1994. Antioxidant actions of thymol, carvacrol, 6-gingerol, zingerone and hydroxytyrosol. *Food and Chemical Toxicology*, 32: 31–36.
- Ahmad I, Mehmood Z, Mohammad F, 1998. Screening of some Indian medicinal plants for their antimicrobial properties. *Journal of Ethnopharmacology*, 62: 183–193.
- Albuquerque MR, Costa SM, Bandeira PN, Santiago GM, Andrade-Neto M, Silveira ER, Pessoa OD, 2007. Nematicidal and larvicidal activities of the essential oils from aerial parts of *Pectis oligocephala* and *Pectis apodocephala* Baker. *Anais da Academia Brasileira de Ciências*. 79: 209–213.
- Ankli A, Heinrich M, Bork P, Wolfram L, Bauerfeind P, Brun R, Schmid C, Weiss C, Bruggisser R, Gertsch J, Wasescha M, Sticher O, 2002. Yucatec Mayan medicinal plants: evaluation based on indigenous uses. *Journal of Ethnopharmacology*, 79: 43–52.
- Ao Y, Satoh K, Shibano K, Kawahito Y, Shioda S, 2008. Singlet oxygen scavenging activity and cytotoxicity of essential oils from rutaceae. *Journal of Clinical Biochemistry and Nutrition*, 43: 6–12.
- Arunasree KM, 2010. Anti-proliferative effects of carvacrol on a human metastatic breast cancer cell line, MDA-MB 231. *Phytomedicine*, 17: 581–588.
- Baik JS, Kim SS, Lee JA, Oh TH, Kim JY, Lee NH, Hyun CG, 2008. Chemical composition and biological activities of essential oils extracted from Korean endemic citrus species. *Journal of Microbiology and Biotechnology*, 18: 74–79.
- Bakkali F, Averbeck S, Averbeck D, Idaomar M, 2008. Biological effects of essential oils – a review. *Food and Chemical Toxicology*, 46: 446–475.
- Baris O, Gulluce M, Sahin F, Ozer H, Kilic H, Ozkan H, Kiliç H, Ozkan H, Sokmen M, Ozbek T, 2006. Biological activities of the essential oil and methanol extract of *Achillea biebersteinii* afan (Asteraceae). *Turkish Journal of Biology*, 30: 65–73.
- Braga PC, Alfieri M, Culici M, Sasso MD, 2007. Inhibitory activity of thymol against the formation and viability of *Candida albicans* hyphae. *Mycoses*, 50: 502-506.
- Chami N, Bennis S, Chami F, 2005. Study of anticandidal activity of carvacrol and eugenol in vitro and in vivo. *Oral Microbiology and Immunology*, 20: 106-111.
- Chiu HF, Wu YC, 2002. Cytotoxic constituents of the stem bark of *Neolitsea acuminatissima*. *Journal of Natural Products*, 65: 255-258.
- Chou TC, Talalay P, 1984. Quantitative analysis of dose-effect relationships: the combined effects of multiple drugs or enzyme inhibitors. *Advances in Enzyme Regulation*, 22: 27-55.
- Clemedson C, Ekwall B, 1999. Overview of the final MEIC results. The in vitro–in vitro evaluation. *Toxicology In Vitro*, 13: 657–663.
- Cragg GM, Newman DJ, 1999. Discovery and development of antineoplastic agents from natural sources. *Cancer Investigation*, 17: 153-163.
- Dutta BK, Rahman I, Das TK, 1998. Antifungal activity of Indian plant extracts. *Mycoses*, 41: 535–536.
- Evangelou A, Kalpousos G, Karkabounas S, Liasko R, Nonni A, Stefanou D, Kallistratos G, 1997. Dose-related preventive and therapeutic effects of antioxidants — anticarcinogens on experimentally induced malignant tumors in Wistar rats. *Cancer Letters*, 115: 105–111.
- Fenaroli G, 2002. *Fenaroli's Handbook of Flavor Ingredients*. CRC Press, Boca Raton, USA.
- He L, Mo H, Hadisusilo S, Qureshi AA, Elson CE, 1997. Isoprenoids suppress the growth of murine B16 melanomas in vitro and in vivo. *Journal of Nutrition*, 127: 668–674.
- Hedberg I, 1993. Botanical methods in ethnopharmacology and the need for conservation of medicinal plants. *Journal of Ethnopharmacology*, 38: 121–128.
- Heinrich M, Gibbons S, 2001. Ethnopharmacology in drug discovery: an analysis of its role and potential contribution. *Journal of Pharmacy and Pharmacology*, 53: 425–432.
- Horvathova E, Turcaniova V, Slamenova D, 2007. Comparative study of DNA damaging and DNA-protective effects of selected components of essential plant oils in human leukemic cells K562. *Neoplasma*, 54: 478–483.
- Iwu MW, Duncan AR, Okunji CO, 1999. New antimicrobials of plant origin. In: Janick J, editor. *Perspectives on new crops and new uses*. Alexandria: ASHS Press; pp. 457–462.

- Kallistratos G, Evangelou A, Agnantis N, Fasske E, Karkabounas S, Donos A, 1994. Enhancement of the antineoplastic effect of anticarcinogens on benzo[a]pyrene-treated Wistar rats, in relation to their number and biological activity. *Cancer Letters*, 82: 153–165.
- Karkabounas S, Binolis J, Zelovitis J, Kotsis N, Charalabopoulos A, Avdikos A, Zouridakis A, Liasko R, Giannakopoulos X, Charalabopoulos K, 2002. Inhibition and modification of benzo[a]pyrene-induced chemical carcinogenesis by ascorbic acid alone or in combination with α -tocopherol in Wistar rats. *Experimental Oncology*, 24: 274–278.
- Karkabounas S, Kostoula OK, Daskalou T, Veltsistas P, Karamouzis M, Zelovitis I, Metsios A, Lekkas P, Evangelou AM, Kotsis N, Skoufos I, 2006. Anticarcinogenic and antiplatelet effects of carvacrol. *Experimental Oncology*, 28: 121–125.
- Katsube N, Iwashita K, Tsushida T, Yamaki K, Kobori M, 2003. Induction of apoptosis in cancer cells by Bilberry (*Vaccinium myrtillus*) and the anthocyanins. *Journal of Agricultural and Food Chemistry*, 51: 68–75.
- Khan A, Ahmad A, Manzoor N, Khan LA, 2010. Antifungal activities of *Ocimum sanctum* essential oil and its lead molecules. *Natural Product Communications*, 5(2): 345–349.
- Koparal AT, Zeytinoglu M, 2003. Effects of carvacrol on a human non-small cell lung cancer (NSCLC) cell line A549. *Cytotechnology*, 43: 149–154.
- Lampronti I, Saab AM, Gambari R, 2006. Antiproliferative activity of essential oils derived from plants belonging to the Magnoliophyta division. *International Journal of Oncology*, 29: 989–995.
- Liasko R, Kabanos T, Karkabounas S, Malamas M, Tasiopoulos A, Stefanou D, Collery P, Evangelou A, 1998. Beneficial effects of a Vanadium complex with cysteine administrated at low doses of benzo[a]pyrene — induced leiomyosarcomas in Wistar rats. *Anticancer Research*, 18: 3609–3613.
- Mastelic J, Jerkovic I, Blazevic I, Poljak-Blazi M, Borovic S, Ivancic-Bace I, Smrecki V, Zarkovic N, Brcic-Kostic K, Vikic-Topic D, Muller N, 2008. Comparative study on the antioxidant and biological activities of carvacrol, thymol, and eugenol derivatives. *Journal of Agricultural and Food Chemistry*, 56: 3989–3996.
- Mukherjee AK, Basu S, Sarkar N, Ghosh AC, 2001. Advances in cancer therapy with plant based natural products. *Current Medicinal Chemistry*, 8: 1467–1486.
- Pina-Vaz C, Rodrigues AG, Pinto E, 2004. Antifungal activity of thymus oils and their major compounds. *Journal of the European Academy of Dermatology and Venereology*, 18: 73–78.
- Pinn G, 2001. Herbs used in obstetrics and gynaecology. *Australian Family Physician*, 30: 351–356.
- Popov AM, Atopkina LN, Uvarova NI, Elyakov GB, 2001. The antimetastatic and immunomodulating activities of ginseng minor glycosides. *Doklady Biochemistry and Biophysics*, 380: 309–312.
- Richardson MA, 2001. Biopharmacologic and herbal therapies for cancer: research update from NCCAM. *Journal of Nutrition*, 131: 3037–3040.
- Ruffa MJ, Ferraro G, Wagner ML, Calcagno ML, Campos RH, Cavallaro L, 2001. Cytotoxic effect of Argentine medicinal plant extracts on human hepatocellular carcinoma cell line. *Journal of Ethnopharmacology*, 79: 335–339.
- Sambrook J, Russell DW, 2001. *Molecular Cloning: A Laboratory Manual*. Third edition. Cold Spring Harbor Laboratory Press, Plainview, NY.
- Scheers ME, Ekwall Ba, Dierickx JP, 2001. In vitro long-term cytotoxicity testing of 27 MEIC chemicals on HepG2 cells and comparison with acute human toxicity data. *Toxicology In Vitro*, 15: 153–161.
- Tatman D, Mo H, 2002. Volatile isoprenoid constituents of fruits, vegetables and herbs cumulatively suppress the proliferation of murine B16 melanoma and human HL-60 leukemia cells. *Cancer Letters*, 175: 129–139.
- Wargovich MJ, Woods C, Hollis DM, Zander ME, 2001. Herbs, cancer prevention and health. *Journal of Nutrition*, 131: 3034–3036.
- Weyermann J, Lochmann D, Zimmer A, 2005. A practical note on the use of cytotoxicity assays. *International Journal of Pharmaceutics*, 288: 369–376.
- Wei Z, Xiaoxia X, Hong C, Jie Z, Xiaobing Z, Ruoyu L, Furong F, 2007. Effect of quercetin on breeding and apoptosis of cervical cancer HeLa cell and on growth of transplanted tumor in nude mice. *Wuhan University Journal of Natural Sciences*, 12 (3): 569–576.